

A PISUM GENE PREVENTING TRANSITION FROM THE VEGETATIVE TO THE REPRODUCTIVE STAGE

Gottschalk, W.

Institute of Genetics, Bonn, Germany

In an M2 family following X-irradiation, some mutant plants arose which were unable to produce flowers (mutant 172 of our collection). The plants showed normal behavior and development with regard to height, growth rate, and physiological activity, but instead of producing inflorescences, they form lateral branches with extremely shortened internodes. The growing points of these lateral branches (and of the main growing point of the stem) remain active, producing leaves and internodes, while the plants of the mother variety are long since mature. Plants remain alive in the field until killed by frost or by heavy mildew attack towards the end of the growing season. When cultivated in the greenhouse, they survived for two years until they succumbed to disease. Surely they would have survived longer under optimal greenhouse conditions without having formed any flowers. We have maintained this mutant type over a period of 11 generations by propagating plants heterozygous for the mutant gene. During this time, with hundreds of plants having been grown, not a single flower was formed.

Histological studies revealed that the growing points of the main stem and the lateral branches remain mitotically active, permanently producing new cell material which differentiates to somatic tissues. Differentiation of floral buds does not occur. Nor did cultivation of the plants in the greenhouse under different photoperiodic conditions induce flower formation. Thus, the behavior of the mutant seems not to be due to a specific photoperiodic requirement. Presumably an enzyme(s) necessary for the transition of the plants from the vegetative to the reproductive stage is lacking.

Mutant 172 arose by X-irradiation. What appears to be the same mutant was selected in neutron experiments. It is like 172 in all respects, but heterozygous plants of the two sources have not yet been crossed to verify their identity. The recessive gene in Mutant 172 is designated "veg".

SIMPLE PROTEIN ASSAY FOR PEA

Hartmann, K. Institute of Genetics, Bonn, West Germany

Breeding for increased protein in field crops has been facilitated by the availability of efficient screening methods. Esen (Anal. Biochem. 89:264-273. 1978) has recently published a simple determination method for quantitative assay of protein which gives a linear response from 0.05 to 3-4 mg/ml of protein. Beside purified proteins like cytochrome C, Bovine serum albumin (BSA), and egg albumin, Esen proposed the applicability of this method for protein extractions of plant tissues. The method requires spotting of five-microliter samples of the protein solutions in **question** on Whatman chromatography paper No. 1. The sheet is stained in a 0.1% coomassie brilliant blue R-250 solution [composed of 65 parts of distilled water, 25 parts of isopropyl alcohol, and 10 parts of acetic acid (v/v/v)] for 15 min. Then the paper is destained by passing it through 3 glass trays, each containing distilled water. Following drying, the stained protein spots are cut out in the form of disks with a corkborer (19 mm diam). An unspotted disk serves as a blank. The dye-protein