

Contents

Review Paper

- Strategies for improving the iron nutritional quality of seed crops: lessons learned from the study of unique iron-hyperaccumulating pea mutants
Michael A. Grusak 1

Research Papers

- A new allele at *Crd* disturbs development of the compound leaf
V.A. Berdnikov, F.L. Gorel', and O.E. Kosterin 6
- Interaction of a new leaf mutation *ins2* with *af*, *uni^{lac}* and *tl^w*
V.A. Berdnikov, F.L. Gorel', V.S. Bogdanova and O.E. Kosterin 9
- New mutations in the linkage group V have provided stable trisomic lines of pea
V.A. Berdnikov, F.L. Gorel' and O.E. Kosterin 13
- A study of potential ability for cross-pollination in pea originating from different parts of the world
V.S. Bogdanova and V.A. Berdnikov 16
- Dominant alleles of the locus *Tl* may differ in their strength
V.S. Bogdanova, O.E. Kosterin and V.A. Berdnikov 18
- Chromosomal location of *Fwf*, the Fusarium wilt race 5 resistance gene in *Pisum sativum*
C.J. Coyne, D.A. Inglis, S.J. Whitehead, M.T. McClendon and F.J. Muehlbauer 20
- Construction of a bacterial artificial chromosome library for pea (*Pisum sativum*L.)
C.J. Coyne, K. Medsem, D.A. Lightfoot, K.E. Keller, R.R. Martin, M.T. McClendon, D.A. Inglis, E.W. Storlie and K.E. McPhee 23
- Selected AFLP primer pairs for fine mapping in pea
C.J. Coyne, M.-L. Pilet, G. Deniot, A. Baranger, S. Prioul and M.T. McClendon..... 27
- Two new field pea cultivars for the southeastern highlands of Ethiopia
T. Getachew..... 31
- Double mutant *rms2 rms5* expresses a transgressive, profuse branching phenotype
I.C. Murfet and G.M. Symons 33
- Location and STS primers for *Cop1*
R.L. Murphy, N.F. Weeden and J.A. Przyborowski 39
- ITS sequence variation in selected taxa of *Pisum*
D.E. Saar and N.O. Polans 42
- Linkages of the *Aba* (*Albumin a*) locus with markers of the linkage group VI
W.K. Swiecicki, J. Przybylska and Z. Zimniak-Przybylska 46
- Densinodosum* – a new gene on linkage group III
W.K. Swiecicki and B. Wolko..... 48
- The position of *Age* and *Wsp* on linkage group IV
N.F. Weeden, M. McBroom, J. Przyborowski and P. Burger 51
- A gene influencing tolerance to common root rot is located on linkage group IV
N.F. Weeden, R. McGee, C.R. Grau and F.J. Muehlbauer 53

Brief Communications

How to identify edible-pod pea varieties in a germplasm collection J.M. Amurrio, A.M. De Ron and J. Hernández-Nistal.....	56
Linkage relationship between the loci <i>Tw1</i> and <i>Vam</i> G.L. Gorel', O.E. Kosterin and V.A. Berdnikov	57
Map position of the <i>Rms6</i> locus I.C. Murfet and C. Rameau	58
The pea <i>rms2-1 rms4-1</i> double-mutant phenotype is transgressive I.C. Murfet and G.M. Symons	59
The position of locus <i>A</i> on the consensus map J. Przyborowshi and N.F. Weeden.....	60
Further confirmatin of the integrity of linkage group III M. Tonguc and N.F. Weeden	62
Four more symbiotic mutants obtained using EMS mutagenesis of line SGE V.E. Tsyganov, V.A. Voroshilova, S.M. Rozov, A.Y. Borisov and I.A.Tikhonovich.....	63

